arXiv:hep-ex/0406045v2 30 Oct 2004

First Observation of a Narrow Charm-Strange Meson $D^+_{sJ}(2632) \to D^+_s\eta$ and D^0K^+

A.V. Evdokimov,[8] U. Akgun,[16] G. Alkhazov,[11] J. Amaro-Reyes,[13] A.G. Atamantchouk,[11,a] A.S. Ayan,[16]
M.Y. Balatz,[8,a] N.F. Bondar,[11] P.S. Cooper,[5] L.J. Dauwe,[17] G.V. Davidenko,[8] U. Dersch,[9,b] A.G. Dolgolenko,[8]
G.B. Dzyubenko,[8] R. Edelstein,[3] L. Emediato,[19] A.M.F. Endler,[4] J. Engelfried,[13,5] I. Eschrich,[9,c]
C.O. Escobar,[19,d] I.S. Filimonov,[10,a] F.G. Garcia,[19,5] M. Gaspero,[18] I. Giller,[12] V.L. Golovtsov,[11] P. Gouffon,[19]
E. Gülmez,[2] He Kangling,[7] M. Iori,[18] S.Y. Jun,[3] M. Kaya,[16,e] J. Kilmer,[5] V.T. Kim,[11] L.M. Kochenda,[11]
I. Konorov,[9,f] A.P. Kozhevnikov,[6] A.G. Krivshich,[11] H. Krüger,[9,g] M.A. Kubantsev,[8] V.P. Kubarovsky,[6]
A.I. Kulyavtsev,[3,5] N.P. Kuropatkin,[11,5] V.F. Kurshetsov,[6] A. Kushnirenko,[3,6] S. Kwan,[5] J. Lach,[5] A. Lamberto,[20]
L.G. Landsberg,[6] I. Larin,[8] E.M. Leikin,[10] Li Yunshan,[7] M. Luksys,[14] T. Lungov,[19] V.P. Maleev,[11] D. Mao,[3,h]
Mao Chensheng,[7] Mao Zhenlin,[7] P. Mathew,[3,i] M. Mattson,[3] V. Matveev,[8] E. McCliment,[16] M.A. Moinester,[12]
V.V. Molchanov,[6] A. Morelos,[13] K.D. Nelson,[16,j] A.V. Nemitkin,[10] P.V. Neoustroev,[11] C. Newsom,[16] A.P. Nilov,[8]
S.B. Nurushev,[6] A. Ocherashvili,[12,k] Y. Onel,[16] E. Ozel,[16] S. Ozkorucuklu,[16,l] A. Penzo,[20] S.V. Petrenko,[6]
P. Pogodin,[16] M. Procario,[3,m] E. Ramberg,[5] G.F. Rappazzo,[20] B.V. Razmyslovich,[11,n] V.I. Rud,[10]
J. Russ,[3] P. Schiavon,[20] J. Simon,[9,o] A.I. Sitnikov,[8] D. Skow,[5] V.J. Smith,[15] M. Srivastava,[19] V. Steiner,[12]
V. Stepanov,[11,n] L. Stutte,[5] M. Svoiski,[11,n] N.K. Terentyev,[11,3] G.P. Thomas,[1] I. Torres,[13] L.N. Uvarov,[11]
A.N. Vasiliev,[6] D.V. Vavilov,[6] E. Vázquez-Jáuregui,[13] V.S. Verebryusov,[8] V.A. Victorov,[6] V.E. Vishnyakov,[8]
A.A. Vorobyov,[11] K. Vorwalter,[9,p] J. You,[3,5] Zhao Wenheng,[7] Zheng Shuchen,[7] and R. Zukanovich-Funchal[19]
(The SELEX Collaboration)

[1]*Ball State University, Muncie, IN 47306, U.S.A.*
[2]*Bogazici University, Bebek 80815 Istanbul, Turkey*
[3]*Carnegie-Mellon University, Pittsburgh, PA 15213, U.S.A.*
[4]*Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*
[5]*Fermi National Accelerator Laboratory, Batavia, IL 60510, U.S.A.*
[6]*Institute for High Energy Physics, Protvino, Russia*
[7]*Institute of High Energy Physics, Beijing, P.R. China*
[8]*Institute of Theoretical and Experimental Physics, Moscow, Russia*
[9]*Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany*
[10]*Moscow State University, Moscow, Russia*
[11]*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
[12]*Tel Aviv University, 69978 Ramat Aviv, Israel*
[13]*Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico*
[14]*Universidade Federal da Paraíba, Paraíba, Brazil*
[15]*University of Bristol, Bristol BS8 1TL, United Kingdom*
[16]*University of Iowa, Iowa City, IA 52242, U.S.A.*
[17]*University of Michigan-Flint, Flint, MI 48502, U.S.A.*
[18]*University of Rome "La Sapienza" and INFN, Rome, Italy*
[19]*University of São Paulo, São Paulo, Brazil*
[20]*University of Trieste and INFN, Trieste, Italy*

We report the first observation of a charm-strange meson $D^+_{sJ}(2632)$ at a mass of 2632.5 ± 1.7 MeV/c^2 in data from SELEX, the charm hadro-production experiment E781 at Fermilab. This state is seen in two decay modes, $D^+_s\eta$ and D^0K^+. In the $D^+_s\eta$ decay mode we observe a peak with 101 events over a combinatoric background of 54.9 events at a mass of 2635.4 ± 3.3 MeV/c^2. There is a corresponding peak of 21 events over a background of 6.9 at 2631.5 ± 2.0 MeV/c^2 in the decay mode D^0K^+. The decay width of this state is < 17 MeV/c^2 at 90% confidence level. The relative branching ratio $\Gamma(D^0K^+)/\Gamma(D^+_s\eta)$ is 0.14 ± 0.06. The mechanism which keeps this state narrow is unclear. Its decay pattern is also unusual, being dominated by the $D^+_s\eta$ decay mode.

PACS numbers: 14.40.Lb, 13.30.Eg, 13.25Ft

In 2003 the BaBar collaboration reported the first observation of a massive, narrow charm-strange meson $D^+_{sJ}(2317)$ below the DK threshold [1]. Confirmation quickly followed from CLEO [2] and BELLE [3]. The CLEO collaboration showed that a higher-lying state, suggested by BaBar, existed and was a partner to the $D^+_{sJ}(2317)$. A number of theory papers suggested different explanations for the unexpectedly low mass of the state, which had been thought to lie above the DK threshold [4, 5, 6, 7, 8, 9]. A prediction relevant to this experiment was that the pattern of parity-doubled states is expected to continue to higher excitations with similar splittings [10].

The SELEX experiment used the Fermilab charged hyperon beam at 600 GeV/c to produce charmed particles in a set of thin foil targets of Cu or diamond. The negative beam composition was approximately half Σ^- and half π^-. The three-stage magnetic spectrometer is shown elsewhere [11, 12]. The most important features are the high-precision, highly redundant, vertex detector that provides an average proper time resolution of 20 fs for charm decays, a 10 m long Ring-Imaging Cerenkov (RICH) detector that separates π from K up to 165 GeV/c [13], and a high-resolution tracking system that has momentum resolution of $\sigma_p/p < 1\%$ for a 150 GeV/c track. Photons are detected in 3 lead glass photon detectors, one following each spectrometer magnet. The photon angular coverage in the center of mass typically exceeds 2π. For this analysis the photon energy threshold was 2 GeV. Previous SELEX $\mathrm{D_s}$ studies showed that most of the signal came from the Σ^- beam [14]. We restrict ourselves in this analysis to the 10×10^9 Σ^--induced interactions.

In this study we began with the SELEX $\mathrm{D_s^\pm} \to \mathrm{K^+K^-}\pi^\pm$ sample used in lifetime and hadroproduction studies [14, 15]. Charged conjugate final states are included here and throughout this paper. The sample-defining cuts are defined in the references. The $\mathrm{D_s}$ meson momentum vector had to point back to the primary vertex with $\chi^2 < 8$ and its decay point must have a vertex separation significance of at least 8 from the primary. Tracks which traversed the RICH ($p \gtrsim 22\,\mathrm{GeV}/c$) were identified as kaons if this hypothesis was most likely. The pion was required to be RICH-identified if it went into its acceptance. There are 544 ± 29 Σ^- induced signal events with these cuts.

Due to high multiplicity, photon detection in an open charm trigger is challenging. SELEX has 3 lead glass calorimeters covering much of the forward solid angle. The energy scale for the detector was set first by using electron beam scans. Then π^0 decays were reconstructed from exclusive trigger data, which selected low-multiplicity radiative final states: $\eta \to \gamma\gamma$ and $\pi^+\pi^-\pi^0$, $\omega \to \pi^+\pi^-\pi^0$ as well as η' and f(1285) mesons [16]. The final energy scale corrections were developed using π^0 decays from the high-multiplicity charm trigger data. Further checks in the charm data set were made using single photon decays, e.g., $\Sigma^0 \to \Lambda\gamma$. The uncertainty in the photon energy scale is less than 2%. Details can be found in Ref. [16].

We selected $\eta \to \gamma\gamma$ candidates in the $\gamma\gamma$ mass range 400-800 MeV/c^2. Each photon of the pair has $E_\gamma > 2$ GeV. The photon pair has $\mathrm{E}_{\gamma\gamma} > 15$ GeV. The $\gamma\gamma$ mass distribution from 10^6 charm-trigger events (0.1% of the data) is shown in Fig. 1 where a η signal over a large combinatoric background is seen. A fit to a Gaussian plus an exponentially falling background yields an η mass of 544.8 ± 2.9 MeV/c^2 consistent with the PDG value [17]. The mass uncertainty for this and all subsequent states is only statistical.



FIG. 1: M($\gamma\gamma$) distribution for photon pairs in the η mass region from 0.1% of the data sample. Results for the fit shown are in Table I. The inset shows the background subtracted η signal. The dark points indicate the η signal region.

The observed resolution is 28 ± 4 MeV/c^2 consistent with the SELEX simulation result, 30 MeV/c^2. The simulation includes all the material in the spectrometer and also reproduces the observed π^0 width as a function of energy [16]. The η signal fraction is 5% within a $\pm$60MeV/c^2 mass window when we eliminate events with more than 5 η candidates in this mass window.

We searched for high-mass charm-strange decays that followed the pattern $\mathrm{D_s}$ plus pseudoscalar meson. We had good acceptance and efficiency for the $\mathrm{D_s}$ η channel. The event selection used included the eta selection above, and the D_s selection described above, which yields a S/N of 4/1. The $\mathrm{D_s}$ momenta are typically 150 GeV/c in the SELEX data set; the $\mathrm{E}_\eta > 15$ GeV energy cut is very loose. We rejected events in which there were more than 5 η candidates in the signal region. This cut removed 18 $\mathrm{D_s}$ candidates (3.3%) while reducing the η candidate list by 20%. The η signal region is shown in Fig. 1. The final sample consisted of 615 η candidates from 526 $\mathrm{D_s}$ candidates.

The results of our search are shown in the M(KK$\pi^\pm$ η) - M(KK$\pi^\pm$) mass difference distribution in Fig. 2(a). In this plot we fixed the η mass at the PDG value [17] by defining an η 4-vector with the measured η momentum and the PDG η mass. A clear peak is seen at a mass difference of 666.9 ± 3.3 MeV/c^2.

To estimate the combinatoric background, we matched each $\mathrm{D_s}$ candidate with η candidates from 25 other sample events to form a event-mixed sample representing the combinatoric background of true single charm production and real η candidates. The event-mixed mass distribu-

Fig.	state	fitted yield	ΔM MeV/c^2	Mass MeV/c^2	Significance $(S-B)/\sqrt{B}$	σ MeV/c^2	Γ MeV/c^2	χ^2/n_d
1	$\eta(548) \to \gamma\gamma$	5087 ± 863		544.8 ± 2.9	13.3 σ	28 ± 4		1.17
2	$D_s^+(2632) \to D_s^+\eta$	43.4 ± 9.1	666.9 ± 3.3	2635.4 ± 3.3	6.2 σ	10.9		1.10
3	$D_s^+(2573) \to D^0K^+$	25 ± 9	705.4 ± 4.3	2569.9 ± 4.3	5.4 σ	4.9	14^{+9}_{-6}	0.77
3	$D_s^+(2632) \to D^0K^+$	13.2 ± 4.9	767.0 ± 2.0	2631.5 ± 2.0	5.3 σ	4.9	$< 17(90\%CL)$	
	$D_s^{*+}(2112) \to D_s^+\gamma$	60.1 ± 14.9	143.7 ± 1.5	2112.0 ± 1.5	4.8 σ	5.9		0.94

TABLE I: Fit results for Figures 1-3 and $D_s^{*+}(2112) \to D_s^+\gamma$.



FIG. 2: M(KK$\pi^\pm$ η) - M(KK$\pi^\pm$) mass difference distribution. Charged conjugates are included. The darker shaded region is the event excess used in the estimation of signal significance. The lighter shaded region is the event-mixed combinatoric background described in the text. The inset shows the difference of the two with a Gaussian fit to the signal. Results for the fit shown are in Table I.

tion was then scaled down by 1/25 to predict the combinatoric background in the signal channel. As can be seen in Fig. 2, the event-mixed background models the background shape very well, but produces no signal peak.

To estimate the signal yield we subtracted the combinatoric background (light shaded area) from the signal data. The resulting difference histogram is plotted in the inset in Fig. 2 in the mass-difference range appropriate to our search ($\mathrm{D}_{\mathrm{sJ}}^+$ masses up to 2900 MeV/c^2). Outside the peak region the data scatter about 0. The width in the $\mathrm{D}^0\mathrm{K}^+$ mode, to be discussed below, is consistent with a 4.9 MeV/c^2 Gaussian. We are insensitive to the natural width in the $\mathrm{D}_{\mathrm{s}}^+\eta$ mode. Therefore we fit the difference histogram with only a Gaussian with no residual background terms. The Gaussian width is fixed at the simulation value of 10.9 MeV/c^2. The fit yield is 43.4 ± 9.1 events at a mass of 2635.4 ± 3.3 MeV/c^2. The reduced χ^2 is 1.10 with a confidence level of 31%.

To assess the Poisson fluctuation probability to observe this excess we note that the resolution is about 1 bin; we take a 6-bin cluster as a signal region and perform a counting experiment. There are 101 events over a background of 54.9 ± 1.5 events giving an excess of 46.1 events in 6 adjacent bins. The Poisson fluctuation probability for this excess is 3×10^{-7} including the uncertainty in the background. A conservative estimate of the fluctuation probability anywhere in the search region (up to 2900 MeV/c^2) is 6×10^{-6}.

The signal does not change with variations of ± 2% in the photon energy scale. We also studied combinations of events in the D_s mass sidebands with η candidates and candidates in the D_s mass peak with events in the η mass sidebands. In all cases only smooth combinatoric backgrounds, as in Fig. 2, were observed. The η population in this sample, corrected for η's from the signal channel, has a signal fraction of 0.12 ± 0.05, 1.4σ higher than the global average of 0.05. It is not possible to separate statistical fluctuation from a possible η enrichment of the overall D_s sample.

A GEANT simulation was also used to determine the overall acceptance for these signals. If we detected the D_s from a $\mathrm{D}_{\mathrm{sJ}}^+(2632) \to \mathrm{D}_{\mathrm{s}}^+\eta$ decay, then 35 ± 2% of the time we also detected the $\eta \to \gamma\gamma$. This acceptance was obtained by embedding $\mathrm{D}_{\mathrm{sJ}}^+(2632) \to \mathrm{D}_{\mathrm{s}}^+\eta$ decay events in existing events from the real data. About 55% of the D_s decays in SELEX come through this high mass state. For comparison, about 25% of the D_s come from D_s (2112) decays and a small fraction from either $\mathrm{D}_{\mathrm{sJ}}^+(2317)$ or $\mathrm{D}_{\mathrm{sJ}}^+(2460)$ decays, which are marginally visible in SELEX data.

The decay $\mathrm{D}_{\mathrm{sJ}}^+(2632) \to \mathrm{D}^0\mathrm{K}^+$ is kinematically allowed. After finding the $\mathrm{D}_{\mathrm{sJ}}^+(2632) \to \mathrm{D}_{\mathrm{s}}^+\eta$ signal we searched for this second decay mode as confirmation. The D^0 sample is the Σ^- induced $D^0 \to K^-\pi^+$ subset of the sample used in our measurement of the D^0 lifetime [18] with tight D^0 cuts ($L/\sigma > 6$, point-back $\chi^2 < 5$, and a good fit to the secondary vertex; $\chi^2/n_d < 3$). The K^+ track is > 46 GeV/c (RICH kaon threshold) and is strongly identified by the RICH as ≥ 10 times more likely to be a kaon than any other hypothesis.

The results are shown in Fig. 3(a) where we see both the known $\mathrm{D}_{\mathrm{sJ}}^+(2573)$ state clearly and another peak above the $\mathrm{D}_{\mathrm{sJ}}^{+}(2573)$. We fit each peak with a Breit-Wigner convolved with a fixed width Gaussian plus a constant background term (as suggested from the wrong-sign data discussed below). The Gaussian resolution is set to the simulation value of 4.9 MeV/c^2. The mass difference and width of the $\mathrm{D}_{\mathrm{sJ}}^{+}(2573)$ returned by the fit, $\Delta M = 705.4 \pm 4.3$ MeV/c^2 and $\Gamma = 14^{+9}_{-6}$ MeV/c^2 , respectively, agree well with the PDG values [17] of $\Delta M = 707.9 \pm 1.5$ MeV/c^2 and $\Gamma = 15^{+5}_{-4}$MeV/c^2. The fitted mass difference of the second Breit-Wigner is 767.0 ± 2.0 MeV/c^2, leading to a mass for the new peak of 2631.5 ± 2.0 MeV/c^2. The fitted yield is 13.2 ± 4.9 events. The signal spread is consistent with the Gaussian resolution, even when plotted in 2.5 MeV/c^2 bins, limiting the possible natural width. For the Breit-Wigner fit we find a limit for the width of < 17 MeV/c^2 at 90% confidence level. This signal has a significance ($[S-B]/\sqrt{B}$) of 5.3 σ in a ±15 MeV/c^2 interval. The mass difference between this signal and the one seen in the $\mathrm{D}_{\mathrm{s}}^{+}\eta$ mode is 3.9 ± 3.8 MeV/c^2 , statistically consistent with being the same mass. Unlike the D_{s} case, the $\mathrm{D}^0\mathrm{K}^+$ decay contributes a small fraction to the SELEX D^0 sample.

Combinatoric background will be equally likely to produce a $\mathrm{D}^0\mathrm{K}^-$ combination (wrong-sign kaon) as a $\mathrm{D}^0\mathrm{K}^+$. The wrong-sign combinations are shown in Fig. 3(b). There is no structure in these data, which fits well to a constant background. We conclude that the peak at 2631.5 MeV/c^2 is real and confirms the observation in the $\mathrm{D}_{\mathrm{s}}^{+}\eta$ mode.

The relative branching ratio $\Gamma(\mathrm{D}^0\mathrm{K}^+)/\Gamma(\mathrm{D}_{\mathrm{s}}^{+}\eta\ )$ must be corrected for the relative acceptances of D^0, D_s, η, and K^+ mesons, for the η, D^0 and D_s^+ branching ratios, the relative acceptances of the $\mathrm{D}_{\mathrm{sJ}}^{+}(2632)$ final states. We estimate the relative acceptance ratio from simulation as 91 ± 3%. The relative branching ratio is $\Gamma(\mathrm{D}^0\mathrm{K}^+)/\Gamma(\mathrm{D}_{\mathrm{s}}^{+}\eta\ ) = 0.14 \pm 0.06$. Relative phase space favors the $\mathrm{D}^0\mathrm{K}^+$mode by a factor of 1.53, making this low branching ratio even more surprising.

In conclusion we combined our clean sample of D_{s} mesons with additional photon pairs which made η candidates to study the $\mathrm{D}_{\mathrm{s}}\eta$ mass spectrum. We observe a clear peak of 43.4 ± 9.1 events with a Poisson fluctuation probability $< 6 \times 10^{-6}$ at a mass difference of 666.9 ± 3.3 MeV/c^2 above the ground state D_{s}. The background shape is well-represented by combinatoric background from event-mixing, as discussed above. A corresponding mass peak is also seen in the $\mathrm{D}^0\mathrm{K}^+$channel with a significance of 5.3 σ at the same mass. The combined measurement of the mass of this state is 2632.5 ± 1.7 MeV/c^2. The SELEX mass scale systematic error is under study, but all charm meson masses measured in SELEX agree with PDG values within 1 MeV/c^2. The state is very narrow, consistent with a width due only to resolution in the $\mathrm{D}^0\mathrm{K}^+$decay mode. The 90% confidence level upper limit for the width is < 17 MeV/c^2.



FIG. 3: (a) D_{s} (2632) → $\mathrm{D}^0\mathrm{K}^+$ mass difference distribution. Charged conjugates are included. The shaded regions are the event excesses used in the estimation of signal significances. Results for the fit shown are in Table I. (b) Wrong sign background D^0 K^- events, as described in the text.

SELEX reports these peaks as the first observation of yet another narrow, high-mass D_{s} state decaying strongly to a ground state charm plus a pseudoscalar meson. The mechanism which keeps this state narrow is unclear. The $\mathrm{D}^0\mathrm{K}^+$ channel is well above threshold, with a Q value ∼ 275 MeV. The branching ratios for this state are also unusual. The $\mathrm{D}_{\mathrm{s}}^{+}\eta$ decay rate dominates the $\mathrm{D}^0\mathrm{K}^+$rate by a factor of ∼ 7 despite having half the phase space.

The authors are indebted to the staff of Fermi National Accelerator Laboratory and for invaluable technical support from the staffs of collaborating institutions. We thank A.M. Badalyan, Bill Bardeen, Estia Eichten, Chris Hill, Maciej Nowak and Yu.A. Simonov for helpful conversations. We gratefully acknowledge helpful suggestions, comments and criticisms from a number of members of the charm community. This project was supported in part by Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie, Consejo Nacional de Ciencia y Tecnología (CONACyT), Conselho Nacional de Desenvolvimento Científico e Tecnológico, Fondo de Apoyo a la Investigación (UASLP), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities, Istituto Nazionale di Fisica Nucleare (INFN), the International Science Foundation (ISF), the National Science Foundation (Phy #9602178),a NATO (grant CR6.941058-1360/94), the Russian Academy of Science, the Russian Ministry of Science and Technology, the Turkish Scientific and Technological Research Board (TÜBİTAK), the U.S. Department of Energy (DOE grant DE-FG02-91ER40664 and DOE contract number DE-AC02-76CHO3000), and the U.S.-Israel Binational Science Foundation (BSF).

Note Added in Proof - At the summer conferences the B-factory experiments[19, 20] and the FOCUS photoproduction experiment[21] all reported negative results in their search for $D_{sJ}(2632)$ production. The SELEX data have an unusually low ratio of $R_s \equiv D_s^{*+}(2112)/D_s^+$ production (Table I) compared to e^+e^- machines. The SELEX R_s ratio can be brought into agreement with the published CLEO result [2] if we *exclude* the D_s events that come through $D_{sJ}(2632)$ decays (55% of the SELEX D_s yield). This fact, together with the strong production asymmetry in the SELEX data, can be interpreted as two-component production of charm-strange states from the Σ^- beam. One component involves normal fragmentation like in photoproduction, the other involves a different mechanism connected with the beam hadron. To understand this production conundrum and to place this new state in the spectroscopy of the charm-strange meson system will require careful study from a number of experiments in the future.

---

[a] Deceased
[b] Present address: Advanced Mask Technology Center, Dresden, Germany
[c] Present address: University of California at Irvine, Irvine, CA 92697, USA
[d] Present address: Instituto de Física da Universidade Estadual de Campinas, UNICAMP, SP, Brazil
[e] Present address: Kafkas University, Kars, Turkey
[f] Present address: Physik-Department, Technische Universität München, 85748 Garching, Germany
[g] Present address: The Boston Consulting Group, München, Germany
[h] Present address: Lucent Technologies, Naperville, IL
[i] Present address: SPSS Inc., Chicago, IL
[j] Present address: University of Alabama at Birmingham, Birmingham, AL 35294
[k] Present address: Sheba Medical Center, Tel-Hashomer, Israel
[l] Present address: Süleyman Demirel Universitesi, Isparta, Turkey
[m] Present address: DOE, Germantown, MD
[n] Present address: Solidum, Ottawa, Ontario, Canada
[o] Present address: Siemens Medizintechnik, Erlangen, Germany
[p] Present address: Allianz Insurance Group IT, München, Germany

[1] B. Aubert et al. (BABAR), Phys. Rev. Lett. **90**, 242001 (2003), hep-ex/0304021.
[2] D. Besson et al. (CLEO), Phys. Rev. **D68**, 032002 (2003), hep-ex/0305100.
[3] P. Krokovny et al. (Belle), Phys. Rev. Lett. **91**, 262002 (2003), hep-ex/0308019.
[4] M. A. Nowak and I. Zahed, Phys. Rev. **D48**, 356 (1993).
[5] M. A. Nowak, M. Rho, and I. Zahed, Phys. Rev. **D48**, 4370 (1993), hep-ph/9209272.
[6] M. A. Nowak, M. Rho, and I. Zahed (2003), hep-ph/0307102.
[7] M. Di Pierro and E. Eichten, Phys. Rev. **D64**, 114004 (2001), hep-ph/0104208.
[8] W. A. Bardeen, E. J. Eichten, and C. T. Hill, Phys. Rev. **D68**, 054024 (2003), hep-ph/0305049.
[9] W. A. Bardeen and C. T. Hill, Phys. Rev. **D49**, 409 (1994), hep-ph/9304265.
[10] W. A. Bardeen, E. J. Eichten, and C. T. Hill, *Private communication.*
[11] M. E. Mattson (SELEX) (2002), FERMILAB-THESIS-2002-03.
[12] J. Russ et al. (SELEX) (1998), hep-ex/9812031.
[13] J. Engelfried et al. (SELEX), Nucl. Instrum. Meth. **A433**, 149 (1999).
[14] M. Kaya et al. (SELEX), Phys. Lett. **B558**, 34 (2003), hep-ex/0302039.
[15] M. Iori et al. (SELEX), Phys. Lett. **B523**, 22 (2001), hep-ex/0106005.
[16] M. Balatz et al. (SELEX), *The lead-glass calorimeter for the selex experiment* (2004), Fermilab/TM/2252(2004).
[17] K. Hagiwara et al. (PDG2002), Phys. Rev. **D66**, 010001 (2002).
[18] A. Kushnirenko et al. (SELEX), Phys. Rev. Lett. **86**, 5243 (2001), hep-ex/0010014.
[19] D. Cronin-Hennessy and R. Galik (CLEO), *Private communication.*
[20] B. Aubert et al. (BABAR) (2004), hep-ex/0408087.
[21] R. Kutschke (FOCUS), *Private communication.*